

## INDEX TO VOLUME V

New names, and the final members of new combinations, are in **bold face type**

- Abies*, 233, 235; *balsamea*, 233-239;  
*pectinata*, 151, 152  
*Abortiporus distortus*, 314  
*Acer Negundo*, 264  
*Acrosporium*, 57, 58, 61; ***compactum***,  
 58; ***Euonymi-japonici***, 58; ***Gos-***  
***sypii***, 59, 61; ***hyalina***, 58; ***leuco-***  
***conium***, 58; *monilioides*, 57, 58;  
*obductum*, 58; ***pirinum***, 58;  
***Tuckeri***, 58  
*Aecia*, Internal, 303  
 Agaricaceae of the Pacific Coast, The  
 —IV. New species of *Clitocybe*  
 and *Melanoleuca*, 206  
 Agaricaceae of tropical North Amer-  
 ica, The—VI, 18  
*Agaricus*, 81; *agglutinatus*, 83; *alu-*  
*taceus*, 310; *asper*, 75; *baccatus*,  
 84; *badius*, 82; *bulbosus*, 74; *bul-*  
*bosus vernus*, 74; *cacaophyllus*, 27;  
*Caesareus*, 73; *campestris*, 72, 169,  
 225; *camphoratus*, 307; *Ceciliae*,  
 86; *chlorinosmus*, 78; *chrysopellus*,  
 23; *chrysotrichus*, 21; *circellatus*,  
 306; *citrinus*, 74; *daucipes*, 84;  
*deliciosus*, 305; *echinocephalus*, 77;  
*excelsus*, 80; *flocculentus*, 36;  
*Frostianus*, 76; *fulvus*, 82; *hel-*  
*voliceps*, 20; *hyalinus*, 82; *inaura-*  
*tus*, 86; *insulsus*, 306; *lactifluus*  
*dulcis*, 307; *Listeri*, 307; *macro-*  
*mastes*, 36; *maculatus*, 80; *magnifi-*  
*cus*, 75; *martianus*, 36; *mitissimus*,  
 307; *monticulosus*, 77; *musaeicola*,  
 27; *muscarius major*, 77; *muscarius*  
*minor*, 76; *myodes*, 75; *nigricans*,  
 308; *nitidus*, 80; *nivalis*, 84; *onus-*  
*tus*, 77; *pantherinus*, 80; *peregrinus*,  
 26; *phalloides*, 74; *piperatus*, 307;  
*plumbeus*, 82; *polypyramis*, 77;  
*porphyrius*, 81; *praetorius*, 85;  
*pustulatus*, 75; *pyrrhus*, 28; *pyrrhus*  
*leiosporus*, 28; *Ravenelii*, 77; *recu-*  
*titus*, 81; *ricensis*, 26; *rubens*, 75;  
*rubescens*, 75; *russuloides*, 77;  
*scobifer*, 35; *scrobiculatus*, 305;  
*soleatus*, 83; *solitarius*, 77; *spissus*,  
 81; *spretus*, 73; *stramineus*, 74;  
*strangulatus*, 85; *strobiliformis*, 77,  
 80; *subdulcis*, 307; *theiogalus*, 306;  
*torminosus*, 305; *trivialis*, 306;  
*vaginatus*, 82; *vellereus*, 307; *ver-*  
*rucusus*, 75; *virosus*, 81; *volvatus*,  
 83; *zonarius*, 306  
*Akebia quinata*, 248  
*Aleuria fulgens*, 302  
 Algal host, The lichen and its. The  
 nature and classification of lichens  
 —II, 97  
*Alternaria*, 279  
*Alysidium*, 47; *fulvum*, 47, 48  
*Amanita*, 72, 226, 229, 230, 232;  
*abrupta*, 79; *Amici*, 77; *ampla*,  
 80; *aspera*, 75; *bisporigera*, 74;  
*Caesarea*, 73, 225, 316; *candida*,  
 78; *cinereoconia*, 78; *citrina alba*,  
 80; *cothurnata*, 74, 96; *crenulata*,  
 77; *elliptosperma*, 79; *elongata*, 79;  
*farinosa*, 84; *flavoconia*, 76; *flavo-*  
*rubens*, 79; *flavorubescens*, 76, 95;  
*floccocephala*, 74; *Frostiana*, 96;  
*glabriceps*, 79; *junquillea*, 77, 80;  
*lignophila*, 74; *livida*, 82; *magni-*  
*velaris*, 80; *Mappa*, 74; *Morrisii*,  
 75; *multisquamosa*, 78; *muscaria*,  
 75, 93, 94, 224-227, 229-232, 316;  
*muscaria coccinea*, 83; *pantherina*,  
 74, 75; ***Peckiana***, 67; *pellucida*,  
 73; *phalloides*, 93, 225, 316;  
*prairicola*, 78; *pubescens*, 85; *radi-*  
*cata*, 78; *recutita*, 74; *rubescens*,  
 75; *spadicea*, 82; *submaculata*, 80;  
*umbrina*, 81; *velatipes*, 75; *verna*,  
 74, 79, 80; *vernalis*, 77; *virosa*, 74  
*Amanitas of eastern North America*,  
 The, 72  
*Amanitopsis*, 81, 84; *adnata*, 84; *ag-*  
*glutinata*, 83; *albocreata*, 84; *bac-*  
*cata*, 83; *farinosa*, 84; *hyperborea*,  
 85; *parcivolvata*, 83; *pubescens*,  
 85; *pulverulenta*, 85; *pusilla*, 83;  
*strangulata*, 85, 86; *vaginata*, 82;  
*volvata*, 83, 94  
 American mycological literature, In-  
 dex to, 41, 91, 182, 251, 284, 317  
*Amerosporium Vanillae*, 40  
*Amsinckia intermedia*, 313  
*Anabena*, 131  
 Analytic work whether colonies of  
 the chestnut blight fungus originate  
 from pycnosporos or ascospores, A  
 method of determining in, 274  
*Anthurus*, 268, 272; *borealis*, 268, 269

## INDEX TO VOLUME V

New names, and the final members of new combinations, are in **bold face type**

- Abies*, 233, 235; *balsamea*, 233-239;  
*pectinata*, 151, 152  
*Abortiporus distortus*, 314  
*Acer Negundo*, 264  
*Acrosporium*, 57, 58, 61; ***compactum***,  
 58; ***Euonymi-japonici***, 58; ***Gos-***  
***sypii***, 59, 61; ***hyalina***, 58; ***leuco-***  
***conium***, 58; *monilioides*, 57, 58;  
*obductum*, 58; ***pirinum***, 58;  
***Tuckeri***, 58  
*Aecia*, Internal, 303  
 Agaricaceae of the Pacific Coast, The  
 —IV. New species of *Clitocybe*  
 and *Melanoleuca*, 206  
 Agaricaceae of tropical North Amer-  
 ica, The—VI, 18  
*Agaricus*, 81; *agglutinatus*, 83; *alu-*  
*taceus*, 310; *asper*, 75; *baccatus*,  
 84; *badius*, 82; *bulbosus*, 74; *bul-*  
*bosus vernus*, 74; *cacaophyllus*, 27;  
*Caesareus*, 73; *campestris*, 72, 169,  
 225; *camphoratus*, 307; *Ceciliae*,  
 86; *chlorinosmus*, 78; *chrysopellus*,  
 23; *chrysotrichus*, 21; *circellatus*,  
 306; *citrinus*, 74; *daucipes*, 84;  
*deliciosus*, 305; *echinocephalus*, 77;  
*excelsus*, 80; *flocculentus*, 36;  
*Frostianus*, 76; *fulvus*, 82; *hel-*  
*voliceps*, 20; *hyalinus*, 82; *inaura-*  
*tus*, 86; *insulsus*, 306; *lactifluus*  
*dulcis*, 307; *Listeri*, 307; *macro-*  
*mastes*, 36; *maculatus*, 80; *magnifi-*  
*cus*, 75; *martianus*, 36; *mitissimus*,  
 307; *monticulosus*, 77; *musaeicola*,  
 27; *muscarius major*, 77; *muscarius*  
*minor*, 76; *myodes*, 75; *nigricans*,  
 308; *nitidus*, 80; *nivalis*, 84; *onus-*  
*tus*, 77; *pantherinus*, 80; *peregrinus*,  
 26; *phalloides*, 74; *piperatus*, 307;  
*plumbeus*, 82; *polypyramis*, 77;  
*porphyrius*, 81; *praetorius*, 85;  
*pustulatus*, 75; *pyrrhus*, 28; *pyrrhus*  
*leiosporus*, 28; *Ravenelii*, 77; *recu-*  
*titus*, 81; *ricensis*, 26; *rubens*, 75;  
*rubescens*, 75; *russuloides*, 77;  
*scobifer*, 35; *scrobiculatus*, 305;  
*soleatus*, 83; *solitarius*, 77; *spissus*,  
 81; *spretus*, 73; *stramineus*, 74;  
*strangulatus*, 85; *strobiliformis*, 77,  
 80; *subdulcis*, 307; *theiogalus*, 306;  
*torminosus*, 305; *trivialis*, 306;  
*vaginatus*, 82; *vellereus*, 307; *ver-*  
*rucusus*, 75; *virosus*, 81; *volvatus*,  
 83; *zonarius*, 306  
*Akebia quinata*, 248  
*Aleuria fulgens*, 302  
 Algal host, The lichen and its. The  
 nature and classification of lichens  
 —II, 97  
*Alternaria*, 279  
*Alysidium*, 47; *fulvum*, 47, 48  
*Amanita*, 72, 226, 229, 230, 232;  
*abrupta*, 79; *Amici*, 77; *ampla*,  
 80; *aspera*, 75; *bisporigera*, 74;  
*Caesarea*, 73, 225, 316; *candida*,  
 78; *cinereoconia*, 78; *citrina alba*,  
 80; *cothurnata*, 74, 96; *crenulata*,  
 77; *elliptosperma*, 79; *elongata*, 79;  
*farinosa*, 84; *flavoconia*, 76; *flavo-*  
*rubens*, 79; *flavorubescens*, 76, 95;  
*floccocephala*, 74; *Frostiana*, 96;  
*glabriceps*, 79; *junquillea*, 77, 80;  
*lignophila*, 74; *livida*, 82; *magni-*  
*velaris*, 80; *Mappa*, 74; *Morrisii*,  
 75; *multisquamosa*, 78; *muscaria*,  
 75, 93, 94, 224-227, 229-232, 316;  
*muscaria coccinea*, 83; *pantherina*,  
 74, 75; ***Peckiana***, 67; *pellucida*,  
 73; *phalloides*, 93, 225, 316;  
*prairicola*, 78; *pubescens*, 85; *radi-*  
*cata*, 78; *recutita*, 74; *rubescens*,  
 75; *spadicea*, 82; *submaculata*, 80;  
*umbrina*, 81; *velatipes*, 75; *verna*,  
 74, 79, 80; *vernalis*, 77; *virosa*, 74  
*Amanitas of eastern North America*,  
 The, 72  
*Amanitopsis*, 81, 84; *adnata*, 84; *ag-*  
*glutinata*, 83; *albocreata*, 84; *bac-*  
*cata*, 83; *farinosa*, 84; *hyperborea*,  
 85; *parcivolvata*, 83; *pubescens*,  
 85; *pulverulenta*, 85; *pusilla*, 83;  
*strangulata*, 85, 86; *vaginata*, 82;  
*volvata*, 83, 94  
 American mycological literature, In-  
 dex to, 41, 91, 182, 251, 284, 317  
*Amerosporium Vanillae*, 40  
*Amsinckia intermedia*, 313  
*Anabena*, 131  
 Analytic work whether colonies of  
 the chestnut blight fungus originate  
 from pycnosporos or ascospores, A  
 method of determining in, 274  
*Anthurus*, 268, 272; *borealis*, 268, 269

- Armillaria mellea*, 38, 314, 316; *mucida*, 38; *subannulata*, 216  
*Arnoldia*, 124, 137  
*Arthonia dispersa*, 139; *punctiformis*, 139; *radiata*, 109, 138; *vulgaris*, 122  
*Arthopyrenia*, 110; *cerasi*, 139; *punctiformis*, 110; *rhyponia*, 139  
 Arthur, J. C., *Uredinales* on *Carex* in North America, 240  
*Ascospores*, A method of determining in analytic work whether colonies of the chestnut blight fungus originate from pycnospores or, 274  
*Aseroe*, 268  
*Aspergillus*, 46, 47  
*Aspidium Thelypteris*, 236, 239  
*Aster*, 242, 264  
*Auriscalpium*, 298  
 Bad year for fleshy fungi, A, 315  
 Banker, H. J., Type studies in the *Hydnaceae*—III. The genus *Sarcodon*, 12; IV. The genus *Phellodon*, 62; V. The genus *Hydnellum*, 194; VI. The genera *Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*, 293  
*Barlaea fulgens*, 302  
*Barya*, 111  
*Basidia*, 105  
*Betula odorata*, 89  
*Biatorina Bouteillii*, 151; *synthesa*, 150  
*Bilimbia*, 105  
*Bjerkandera*, 313  
*Blight fungus* originate from pycnospores or ascospores, A method of determining in analytic work whether colonies of the chestnut, 274  
*Boletus*, 176; *americanus*, 3; *flavidus*, 4  
*Botrydina vulgaris*, 114  
*Botrytis*, 46  
*Buellia parasema*, 148; *punctiformis*, 122, 123  
 Burlingham, G. S., The *Lactarieae* of the Pacific Coast, 305  
*Byssonectria*, 177  
*Caeoma Abietis-canadensis*, 238; *nitens*, 281, 282  
*Calicium*, 105; *curtum*, 123; *parietinum*, 123; *trachelinum*, 123  
*Caloscypha*, 299  
*Calospora Vanillae*, 39  
*Calosypha fulgens*, 302  
*Calvatia cyathiformis*, 316; *maxima*, 316  
*Cantharellus*, 261, 262; *brevipes*, 261, 262; *clavatus*, 261, 263; *floccosus*, 262  
*Cantharellus brevipes* and *Cantharellus clavatus*, The identity of, 261  
*Cantharellus clavatus*, The identity of  
*Cantharellus brevipes* and, 261  
*Carex*, 240-244  
*Carex* in North America, *Uredinales* on, 240  
*Catillaria denigrata*, 123; *prasina*, 123  
*Cephaleuros Henningsii*, 40  
*Ceriumyces auriporus*, 2; *bicolor*, 4; *communis*, 260; *fumosipes*, 259; *illudens*, 260; *Peckii*, 5; *speciosus*, 5; *subglabripes*, 4  
*Chaenotheca*, 113; *chrysocephala*, 113  
*Chalara mycoderma*, 45  
*Chamonixia*, 313  
*Chanterel cinnabarinus*, 258; *minor*, 257  
*Chestnut blight fungus* originate from pycnospores or ascospores, A method of determining in analytic work whether colonies of the, 274  
*Chionanthus virginica*, 248  
*Chlorella*, 129; *viridis*, 131  
*Chlorococcum*, 119, 125, 127, 145, 146; *humicola*, 109, 115, 117, 118, 124, 125, 129-131  
*Chondromyces aurantiacum*, 60  
*Chromocrea*, 179  
*Chromocroepsis*, 179  
*Chroococcus*, 126, 131  
*Cladonia*, 131, 150; *pyxidata*, 137  
*Cladotrix*, 125  
 Clark, E. D., & Smith, C. S., Toxicological studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*, 224  
 Classification of lichens, The nature and—II. The lichen and its algal host, 97  
*Clathrella chrysomycelina*, 268; *Clathrus*, 313  
*Clathrogaster*, 313  
*Clathrus*, 267, 268, 271; *cancellatus*, 267  
*Clavaria mucida*, 115; *pistillaris*, 262, 263  
*Claviceps*, 178  
*Climacodon*, 293  
*Clitocybe*, 207, 210, 225, 226, 229-231; *albicastranea*, 206; *albiformis*, 206; *atrialba*, 207, 208; *avellanefalba*, 207; *brunnescens*, 208; *cuticolor*, 208; *cyathiformis*, 208; *dealbata sudorifica*, 225; *griseifolia*, 208; *Harperi*, 209; *hondensis*, 209; *illudens*, 225-227, 229-232; *multiceps*, 225, 230, 232; *murinifolia*, 210; *oculata*, 210; *oreades*, 210; *oregonensis*, 211; *Peckii*, 211; *sinopica*, 212; *stipitata*, 211; *subcandicans*, 212; *subfumosipes*, 212; *subinversa*, 212; *variabilis*, 213; *variabilis*

- brevipes**, 213; **violaceifolia**, 213; **washingtonensis**, 214  
*Clitocybe* and *Melanoleuca*, New species of. The Agaricaceae of the Pacific Coast—IV, 206  
*Clitocybe illudens* and *Inocybe infida*, Toxicological studies on the mushrooms, 224  
*Clitopilus*, 314; **abortivus**, 314  
Coast, The Agaricaceae of the Pacific—IV. New species of *Clitocybe* and *Melanoleuca*, 206  
Coast, The Lactariaceae of the Pacific, 305  
*Coccoloba uvifera*, 246, 247  
*Cocomyxa subellipsoidea*, 114  
*Collema*, 112, 115, 119, 120, 124; **microphyllum**, 123; **pulposum**, 107, 130  
*Colletotrichum*, 171  
*Collybia*, 216, 217; **platyphylla**, 207, 208; **subdecumbens**, 68; **truncata**, 68  
Colonies of the chestnut blight fungus originate from pycnosporos or ascospores, A method of determining in analytic work whether, 274  
Color guide, Ridgway's new, 172  
*Colus* **Garciae**, 268; **hirudinosus**, 268  
Conrad, H. S., The structure of *Simblum sphaerocephalum*, 264  
*Conida*, 114; **punctatella**, 114; **rubescens**, 114  
*Coniophyrium Chionanthi*, 248  
*Conocybe tener*, 36  
*Cookeina*, 185-187; **Afzelii**, 189; **Colensoi**, 185, 187, 191, 193; **Hindsii**, 189; **insititia**, 185, 187, 190, 193; **sulcipes**, 187, 189, 190, 192, 193; **Tricholoma**, 186-188, 193  
*Coprinus*, 168; **atramentarius**, 168, 316; **comatus**, 168, 315  
*Cora*, 115, 125, 126  
*Cordyceps*, 178, 179; **Cockerellii**, 179  
*Coriolus*, 313; **prolificans**, 288; **versicolor**, 287  
*Corticium salmonicolor*, 39  
*Craterellus*, 261, 263; **clavatus**, 261-263; **corrugis**, 263; **pistillaris**, 262, 263  
*Creolophus*, 293; **agaricoides**, 294; **pulcherrimus**, 294; **septentrionalis**, 293  
*Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*, The genera. Type studies in the Hydnaceae—VI, 293  
*Crepidotus*, 18, 26; **alveolus**, 27, 31, 32; **aquosus**, 30; **bicolor**, 28; **cacaophyllus**, 27; **calolepidoides**, 30; **calolepis**, 31; **cinchonensis**, 30; **Citri**, 27, 30; **croceosanguineus**, 28; **cuneiformis**, 29; **Dussii**, 28; **fumosifolius**, 31; **laceratus**, 29; **mollis**, 32; **musaccola**, 27, 32; **parvulus**, 27; **Psychotriace**, 27; **pyrrhus**, 28, 29; **subcuneiformis**, 29; **substipitatus**, 31; **sulcatus**, 29, 30  
*Cryptodiscus araneo-cinctus*, 248  
*Cryptoporus*, 313  
Cultures of heteroecious rusts, Furrer, 233  
*Cup-fungi*, Some tropical, 185  
*Cylindrium*, 45, 46, 55  
*Cypheium*, 118  
*Cystococcus*, 129, 130, 145; **humicola**, 145, 146  
Cytology of the Laboulbeniales, Faull's, 174  
*Dactylococcus*, 125; **infusionum**, 118  
*Daedalea*, 313; **quercina**, 115, 291  
*Dendrogaster*, 313  
*Diaporthe Batatatis*, 279; **parasitica**, 90  
*Diatrype patella*, 249  
*Dicaeoma*, 240-243  
*Dictyonema*, 115, 118, 126  
*Dictyophora irpicina*, 269  
*Didymascella*, 7, 8; **Oxycedri**, 8, 9  
*Didymium anomalum*, 250  
*Diplodia Akebiae*, 248  
*Diploschistes*, 129  
Dodge, B. O., Faull's cytology of the Laboulbeniales, 174  
Durand, E. J., The genus *Keithia*, 6  
Eastern North America, The *Amanitas* of, 72  
*Echinodontium*, 295, 296; **tinctorium**, 295  
*Echinodontium*, *Gloiodon*, and *Hydnodon*, The genera *Creolophus*. Type studies in the Hydnaceae—VI, 293  
*Elfvigia megaloma*, 291  
*Endocarpon pusillum*, 122, 123  
*Endomyces scytonematum*, 113  
*Endophyllum*, 281, 282  
*Endothia*, 274; **gyrosa**, 280; **parasitica**, 90, 274, 280; **radicalis**, 90, 280; **radicalis mississippiensis**, 280  
*Enerthenema syncarpon*, 250  
*Entoloma*, 259; **Grayanum**, 259; **lividum**, 259; **mirabile**, 68; **sinuatum**, 259  
*Ephebe*, 112, 116, 124; **Hegetschweileri**, 113; **pubescens**, 109, 119  
*Epigloea bactrospora*, 111, 118, 120  
*Erigeron*, 242  
*Erysiphe graminis*, 58  
*Eucalyptus*, 35  
*Euglena viridis*, 131

- Eusynchytrium*, 315  
*Evernia furfuracea*, 130; *prunastri*, 149  
*Exobasidium*, 88, 89  
*Exosporium Ulmi*, 40  
  
*Fagus*, 245  
 Fairman, C. E., Notes on new species of fungi from various localities, 245  
 Faull's cytology of the Laboulbeniales, 174  
*Favolus*, 313  
 Fink, Bruce, The nature and classification of lichens—II. The lichen and its algal host, 97  
*Flammula*, 18; **areolata**, 36; **aureobrunnea**, 19; **aureoviridis**, 19; **bryophila**, 36; **chrysotrichoides**, 36; **depressa**, 36; **Earlei**, 36; **hispidula**, 24; **hispidella**, 36; **hypholomoides**, 36; **jalapensis**, 36; **lateritia**, 19; **Nashii**, 36; **olivacea**, 18; **palmicola**, 36; **parvula**, 36; **pholiotoides**, 36; **subpenetrans**, 36; **tenuis**, 36; **vinicolor**, 18  
 Fleshy fungi, A bad year for, 315  
*Fomes*, 295, 296, 313; *Laricis*, 287; *semitostus*, 39; *tinctorius*, 295; *ungulatus*, 291  
 Fraser, W. P., Further cultures of heteroecious rusts, 233  
*Fraxinus*, 247  
*Fuligo megaspora*, 250  
 Fungi, A bad year for fleshy, 315  
 Fungi from various localities, Notes on new species of, 245  
 Fungi, Illustrations of—XIII, 1; XIV, 93; XV, 257; XVI, 287  
 Fungi, New species of, 67  
 Fungi, Some tropical cup-, 185  
 Fungus originate from pycnosporos or ascospores, A method of determining in analytic work whether colonies of the chestnut blight, 274  
 Further cultures of heteroecious rusts, 233  
*Fusarium*, 178, 180, 181, 279; *Solani*, 181; *vasinfectum*, 181  
 Fusarium problem, Wollenweber's studies on the, 178  
*Fusicladium Vanillae*, 40  
*Fusidium*, 45, 46  
  
*Galera frustulenta*, 36  
*Ganoderma*, 313  
*Gautieria*, 313  
*Gaylussacia baccata*, 238  
 Genera *Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*. Type studies in the Hydnaceae—VI, 293  
 Genus *Hydnellum*, The. Type studies in the Hydnaceae—V, 194  
 Genus *Keithia*, The, 6  
 Genus *Phellodon*, The. Type studies in the Hydnaceae—IV, 62  
 Genus *Pseudoplectania*, The, 299  
 Genus *Sarcodon*, The. Type studies in the Hydnaceae—III, 12  
 Genus *Synchytrium*, The, 315  
*Geopyxis aluticolor*, 191; *elata*, 189, 190; *Mölderiana*, 192  
*Geotrichum*, 45, 56; *candidum*, 56, 61; **cuboideum**, 56, 61  
*Gloeocapsa*, 111, 124, 134; *polydermatica*, 118  
*Gloeocystis*, 112  
*Gloeopeziza Rehmii*, 112  
*Gloeosporium*, 171; *Caryae*, 88  
*Gloiodon*, 296, 298  
*Gloiodon*, and *Hydnodon*, The genera *Creolophus*, *Echinodontium*. Type studies in the Hydnaceae—VI, 293  
*Glomerella*, 171; *cingulata*, 171; *Gossypii*, 171; *lindemuthianum*, 171  
*Glomerella*, Shear's studies of parasitic species of, 171  
*Glycophila*, 45, 46, 57  
*Gnomonia Caryae*, 88; *setacea macrospora*, 88  
*Gossypium*, 59  
*Graphis*, 105; *scripta*, 122, 123, 139  
*Grifola frondosa*, 290  
 Guide, Ridgway's new color, 172  
*Gymnoglossum*, 313  
*Gymnophilus*, 18, 21, 35, 36; **areolatus**, 24, 36; **aureobrunneus**, 19; **aureoviridis**, 19; **bryophilus**, 22, 36; *carbonarius*, 25; **chrysopellus**, 20, 23; **chrysotrichoides**, 21, 36; **chrysotrichus**, 21; **depressus**, 20, 36; **Earlei**, 22, 36; **helvoliceps**, 20; **hispidellus**, 24, 36; **hispidus**, 24; **hypholomoides**, 26, 36; **jalapensis**, 25, 36; **lateritius**, 19, 23; **Nashii**, 23, 36; **olivaceus**, 18; **palmicola**, 23, 36; **parvulus**, 19, 36; *penetrans*, 20, 26, 36; **pholiotoides**, 24, 36; *sapineus*, 26; **subpenetrans**, 20, 36; **tenuis**, 22, 36; **vinicolor**, 18  
*Gymnosporangium*, 278  
*Gyroporus castaneus*, 1  
  
*Haematococcus pluvialis*, 131  
*Haematomma ventosum*, 143  
*Halobysus*, 45, 46  
*Haplochytrium*, 315  
 Harper, E. T., The identity of *Cantharellus brevipes* and *Cantharellus clavatus*, 261; The probable identity of *Stropharia epimyces* (Peck) Atk. with *Pilosace algeriensis* Fries, 167

- Heald, F. D., A method of determining in analytic work whether colonies of the chestnut blight fungus originate from pycnospores or ascospores, 274
- Helicocephalum, 45, 46
- Helminthosporium, 279
- Helotium purpuratum, 192
- Hendersonia **coccolobina**, 247; **hypocarpa**, 246; Opuntiae, 38; Rosae, 247
- Heppia urceolata, 124
- Herpotrichia, 282; nigra, 282, 283
- Heteroecious rusts, Further cultures of, 233
- Heuchera cylindrica, 71
- Host, The lichen and its algal. The nature and classification of lichens—II, 97
- Hyalopora, 237
- Hydnaceae, Type studies in the—III. The genus Sarcodon, 12; IV. The genus Phellodon, 62; V. The genus Hydnellum, 194; VI. The genera Creolophus, Echinodontium, Gloiodon, and Hydnodon, 293
- Hydnellum, 65, 198, 199, 204; **Dia-bolus**, 194; **geogenium**, 204; **hybridum**, 198; **inguinatum**, 202; **parvum**, 200; **Peckii**, 203; **Rickerii**, 201; **sanguinarium**, 196, 198; **scrobiculatum**, 196; **suaveolens**, 201, 202; **velutinum**, 196; **Vespertilio**, 198; **zonatum**, 199, 201
- Hydnellum, The genus. Type studies in the Hydnaceae—V, 194
- Hydnodon**, 297; **thelephorum**, 297
- Hydnodon, The genera Creolophus, Echinodontium, Gloiodon and. Type studies in the Hydnaceae—VI, 293
- Hydnofomes, 295, 296; **tsugicola**, 295
- Hydnophysa, 295, 296
- Hydnum, 288, 296, 298; **acre**, 13; **agaricoides**, 294; **amicum**, 62; **boreale**, 201; **bubalinum**, 14; **carbunculus**, 194, 195; **cervinum**, 14, 15; **compactum**, 201; **concresecens**, 199, 200; **corrugatum**, 293; **cristatum**, 13; **cyathiforme**, 64, 65, 199, 200; **discolor**, 294; **ferrugineum**, 195–198; **fragile**, 12; **fragrans**, 64; **friabile**, 204, 295; **fuligineo-violaceum**, 14; **geogenium**, 204, 205; **graveolens**, 62–64; **hybridum**, 197–199; **imbricatum**, 12, 14, 15; **laevigatum**, 14; **leptopus**, 62, 63; **leptopus graveolens**, 64; **lateritium**, 297, 298; **melaleucum**, 62–65; **nigrum**, 62, 198; **Queletii**, 198; **parasiticum**, 297; **pulcherrimum**, 294, 295; **pullum**, 62, 63; **scrobiculatum**, 196–198; **septentrionale**, 293; **spongiosipes**, 196; **stratosum**, 297; **strigosum**, 297; **suaveolens**, 201; **sulphureum**, 204, 205; **thelephorum**, 297; **tomentosum**, 64, 65; **vellereum**, 62; **velutinum**, 196, 197; **Vespertilio**, 198; **zonatum**, 62, 198–200
- Hymenochaete noxia, 39
- Hypheothrix Zenkeri, 112
- Hypophoma, 31; **sublateritium**, 26
- Hyphomycetes, Studies in North American—II, 45
- Hypocrea, 179
- Hypocreaceae, Maire's remarks on some, 176
- Hypodendrum, 18, 35; **scobifer**, 35
- Hypomyces, 176, 179, 180; **armeniacus**, 177; **aureo-nitens**, 177; **boletinus**, 176; **chrysospermus**, 176; **hyalinus**, 177; **inaequalis**, 177; **macrosporus**, 177; **ochraceus**, 177; **papyraceus**, 177; **Solani**, 181; **tegillum**, 177; **violaceus**, 177
- Hypophyllum album, 308
- Hysterangium, 313
- Hysterium, 105
- Identity of Cantharellus brevipes and Cantharellus clavatus, The, 261
- Identity of Stropharia epimyces (Peck) Atk. with Pilosace algeriensis Fries. The probable, 167
- Illustrations of fungi—XIII, 1; XIV, 93; XV, 257; XVI, 287
- Index to American mycological literature, 41, 91, 182, 251, 284, 317
- Inocybe, 224–226, 228, 229, 231; **decipiens**, 225; **infelix**, 224, 225; **infida**, 224–232; **minima**, 69
- Inocybe infida, Toxicological studies on the mushrooms Clitocybe illudens and, 224
- Inonotus hirsutus, 297
- Internal aecia, 303
- Irpex, 288
- Irpiciporus mollis, 288
- Ischnoderma, 313
- Jungermannia, 111; **triophylla**, 112
- Juniperus, 6–8, 11; **communis**, 8; **Oxycedrus**, 9
- Kalchbrennera, 268
- Karschia, 113, 114; **destructans**, 113
- Keithia, 6–8; **tetraspora**, 6–11; **thujina**, 6–9, 11; **Tsugae**, 7, 8, 10, 11
- Keithia, The genus, 6
- Laboulbenia chaetophora, 174; **Gyrindarum**, 174
- Laboulbeniales, Faull's cytology of the, 174
- Lachnea crispata, 192; **nigrella**, 301

- Lactaria, 305; camphorata, 307; Chelidonium, 305; circellata, 307; deliciosa, 305; grisea, 306; insulsa, 306; mitissima, 307; mucida, 306; piperata, 307; scrobiculata, 305; subdulcis, 307; theiogala, 306; torminosa, 305; trivialis, 306; vellerea, 307, 308; zonaria, 306
- Lactariceae of the Pacific Coast, The, 305
- Lactarius brevipes, 306; brevis, 306; deflexus, 306; exsuccus, 308; subserifluus, 307; villosus, 305; xanthogalactus, 307
- Laterna, 268
- Laudatea, 115, 126
- Leaia, 206
- Lecanora atrorufa, 127; caesiorufa, 127; coilocarpa, 127; ferruginea, 127; granatina, 118; pallida, 140; subfusca, 122, 125, 127
- Lecidea, 105; atrobrunnea, 140; enteroleuca, 122
- Lepiota, 94; cretacea, 85; procera, 257
- Leptogonium subtile, 119
- Leptonia gracilipes, 69, 70; validipes, 70
- Leptospora Musae, 89
- Lichen and its algal host, The. The nature and classification of lichens—II, 97
- Lichens, The nature and classification of—II. The lichen and its algal host, 97
- Lichina, 138; confinis, 124; pygmaea, 119
- Liquidambar, 295
- Liriodendron, 54
- Literature, Index to American mycological, 41, 91, 182, 251, 284, 317
- Lycoperdon giganteum, 52
- Lycopus virginicus, 303, 304
- Lysurus, 268
- Maire's remarks on some Hypocreaceae, 176
- Malbranchea, 45, 57; pulchella, 57; pulveracea, 57, 61
- Mangifera indica, 247
- Marasmius, 249
- Melampsora arctica, 238, 239; Medusae, 238, 239
- Melanoleuca, 206; anomala, 214, 223; arenicola, 214, 223; avellanea, 215, 223; avellaneifolia, 215, 223; bicolor, 215, 223; californica, 216, 223; collybiiformis, 216, 223; dryophila, 217, 220, 223; farinacea, 217, 223; Harperi, 217, 223; nuciolens, 218, 223; Olesonii, 218, 223; oreades, 218, 223; pinicola, 219, 223; platyphylla, 219, 223; portolensis, 219, 223; roseibrunnea, 216, 220, 223; rudericola, 220, 223; secedifolia, 221, 223; striatella, 221, 223; subulurida, 221, 223; submulticeps, 221, 223; subpessundata, 217, 222, 223; subvelata, 222, 223; tenuipes, 223
- Melanoleuca, New species of Clitocybe and. The Agaricaceae of the Pacific Coast—IV, 206
- Melascypha, 299; melaena, 300
- Merulius clavatus, 263; lacrymans, 89
- Method of determining in analytic work whether colonies of the chestnut blight fungus originate from pycnosporos or ascospores, A, 274
- Michelia fuscata, 120
- Microsphaera Platani, 58
- Monilia, 45-47; Aspergillus, 47; aurantiaca, 48; aurea, 47; aureofulva, 48; Avenae, 59; candida, 47, 52, 60; capitata, 47; Cerasi, 51; cespitosa, 47; cespitosa aurea, 47; cinerea, 50; crustacea, 47; diffusa, 60; divaricata, 47; effusa, 48; fructigena, 50; fungicola, 51; fusconigra, 60; glauca, 47; globosa, 60; Harknessii, 59; hyalina, 57, 58; Linhartiana, 51, 60; Martinii, 52; megalospora, 49; mycophila, 60; nidulans, 47; Peckiana, 52, 60; penicillata, 60; pulveracea, 57; punctans, 60; racemosa, 47; ramosa, 47; rosea, 47; rubiginosa, 60; simplex, 47; sitophila, 52; urediniformis, 60; viridi-flava, 60
- Mucor Aspergillus, 47; crustaceus, 47
- Murrill, W. A., A bad year for fleshy fungi, 316; Illustrations of fungi—XIII, 1; XIV, 93; XV, 257; XVI, 287; Sterility in Pholiota candicans (Bull.) Schroet., 314; The Agaricaceae of the Pacific Coast—IV. New species of Clitocybe and Melanoleuca, 206; The Agaricaceae of tropical North America—IV, 18; The Amanitas of eastern North America, 72
- Mushrooms Clitocybe illudens and Inocybe infida, Toxicological studies on the, 224
- Mycena, 36
- Mycobacidia, 105
- Mycobilimbia, 105
- Mycocalicium, 105
- Mycological literature, Index to American, 41, 91, 182, 251, 284, 317
- Mycosphaerella lageniformis, 249
- Naevia canadica, 249



- Nature and classification of lichens, The—II. The lichen and its algal host, 97
- Nectandra, 22
- Nectria, 112, 179, 180; *Ipomoeae*, 180; *phytophila*, 112; *Solani*, 181; *Vanillae*, 40
- Nectriopsis, 176, 177, 179
- Neocosmospora, 181; *vasinfecta*, 181
- Neopectia, 282; *Coulteri*, 282, 283
- New color guide, Ridgway's, 172
- New species of *Clitocybe* and *Melanoleuca*. The *Agaricaceae* of the Pacific Coast—IV, 206
- New species of fungi, 67
- New species of fungi from various localities, Notes on, 245
- News and notes, 37, 87, 249, 278, 312; and reviews, 170
- Nigredo, 240–242
- North America, The *Agaricaceae* of tropical—IV, 18
- North America, The *Amanitas* of eastern, 72
- North America, *Uredinales* on *Carex* in, 240
- North American *Hyphomycetes*, Studies in—II, 45
- Nostoc, 115, 119, 120, 124, 125, 130, 134, 136–138; *lichenoides*, 118
- Notes, News and, 37, 87, 249, 278, 312; and reviews, 170
- Notes on new species of fungi from various localities, 245
- Ocellaria *Vanillae*, 40
- Oidium, 45, 47, 48; *albipes*, 59; *Asteris-punicel*, 59; *aureum*, 48, 61; *candidum*, 59; *compactum*, 58; *corticale*, 60; *erysiphoides*, 59; *Euonymi-japonici*, 58; *fructigenum*, 50; *inquinans*, 60; *irregulare*, 60; *Lactis*, 52, 53; *leucoconium*, 58; *leucogonium*, 58; *leuconium*, 58; *megalosporum*, 48, 49, 61; *Murrilliae*, 48, 49, 61; *obductum*, 58; *pirinum*, 58; *simile*, 48, 61; *Tuckeri*, 58
- Ombrophila *limosa*, 249
- Omphalaria, 124
- Onoclea, 237; *sensibilis*, 234, 236, 237, 239; *Struthiopteris*, 234–236, 239
- Oospora, 45, 49, 53, 61; *Arthuri*, 50, 52, 61; *candida*, 50, 52; *candidula*, 55; *Cerasi*, 50, 51, 61; *cinerea*, 50, 61; *crustacea*, 45; *cuboidea*, 56; *cucumeris*, 60; *fructigena*, 50, 61; *fungicola*, 50, 51, 61; *heterospora*, 60; *hyalinula*, 55; *Lactis*, 45, 53; *Linhartiana*, 50–52, 61; *Martinii*, 50, 52, 61; *Nicotianae*, 54; *pallida*, 60; *scabies*, 61; *similis*, 48; *Tulipiferae*, 54
- Oosporoidea, 52; *Lactis*, 53
- Opegrapha, 120, 124; *filicina*, 124; *subsiderella*, 122, 123; *varia*, 119, 120, 124; *vulgata*, 127
- Ophionectria, 179
- Oplismenus *hirtellus*, 246
- Oscillatoria, 125
- Osmunda, 235; *Claytoniana*, 234, 235, 239
- Otidea *domingensis*, 192
- Otidella, 299; *fulgens*, 302; *nigrella*, 301
- Ovularia, 61; *isarioides*, 61
- Pacific Coast, The *Agaricaceae* of the —IV. New species of *Clitocybe* and *Melanoleuca*, 206
- Pacific Coast, The *Lactariae* of the, 305
- Paepalopsis, 45, 46
- Palmella, 111, 112; *botryoides*, 111, 117, 118, 120
- Panaeolus *epimyces*, 167, 168
- Pannaria *brunnea*, 138; *muscorum*, 124; *nigra*, 120; *tryptophylla*, 118, 124
- Parasitic species of *Glomerella*, Shear's studies of, 171
- Parmelia *acetabulum*, 127, 129; *aspidota*, 133; *olivacea*, 152; *physodes*, 148
- Parüphrädria, 111; *Heimerlii*, 111
- Patinella, 105
- Peck, C. H., New species of fungi, 67
- Peckia, 176
- Peltandra *virginica*, 304
- Peltigera, 131; *canina*, 130
- Penicillium, 47
- Peridermium, 235; *balsameum*, 233–237; *Peckii*, 11, 238
- Perisporium *Wrightii*, 38
- Pertusaria, 129; *communis*, 122, 125, 129
- Pestalotia, 245; *truncata* Rubi, 245; *truncata septoriana*, 245
- Pezicula *eximia*, 249
- Peziza, 187; *Afzelii*, 189; *aluticolor*, 191; *Colensoi*, 191; *crispata*, 192; *cyanoderma*, 302; *domingensis*, 186, 192; *fulgens*, 302; *fuscocana*, 300, 301; *Harmoge*, 192; *Hindsii*, 186, 188, 189; *Hystrix*, 186, 188; *insititia*, 190; *melana*, 300; *melania*, 300, 302; *nigrella*, 299, 301; *onotica*, 186; *spongiosa*, 300, 302; *strispora*, 186, 188, 189; *stygia*, 301; *sulcipes*, 188–190; *Tricholoma*, 187–189; *vogesiana*, 300, 302
- Phacidium *tetrasporum*, 6, 8
- Phaeolus, 313
- Phallo-gaster, 313
- Phegopteris *Dryopteris*, 234–236, 239



- Phellinus, 313  
 Phellodon, 65, 66, 298; **amicus**, 62-64; **carnosus**, 65; **niger**, 62; **pul-lus**, 62-64; **tomentosus**, 64  
 Phellodon, The genus. Type studies in the Hydnaceae—IV, 62  
 Phillipsia, 186, 192; **domingensis**, 192, 193; **kermesina**, 192; **subpurpurea**, 192  
 Pholiota, 18, 21, 25, 32, 35, 39; **avellanea**, 32; **Brittoniae**, 35; **Broadwayi**, 32; **bryophila**, 33; **cinchonensis**, 33; **cubensis**, 34; **martinicensis**, 34; **Musae**, 34; **praecox**, 314; **unicolor**, 33; **ventri-cosa**, 35  
 Pholiota candicans (Bull.) Schroet., Sterility in, 314  
 Pholiotina Musae, 34  
 Phoma Batatae, 279; **pigmentivora**, 40; **rhodocarpa**, 246  
 Phyllactidium, 124, 125, 136  
 Phyllosticta **Mortoni**, 247; **Vanillae**, 40  
 Physalis, 264  
 Physcia, 130; **apiolia**, 137; **ciliaris**, 125; **pulverulenta**, 125, 131; **stel-laris**, 127, 148, 152  
 Physma, 137; **chalazanum**, 124  
 Pilocratera, 185, 187; **Engleriana**, 189, 190; **Hindsii**, 189; **Tricholoma**, 188  
 Pilosace algeriensis, 167-169  
 Pilosace algeriensis Fries, The prob-able identity of Stropharia epimyces (Peck) Atk. with, 167  
 Pinus Strobis, 67  
 Piptoporus, 313  
 Pirus coronaria, 58  
 Placodium murorum, 129, 132  
 Placynthium, 119  
 Platanus orientalis, 59, 61  
 Plectania nigrella, 301  
 Pleurococcus, 125, 127, 129, 130, 145; **vulgaris**, 118, 130, 131  
 Pleurotus ulmarius, 316  
 Poa pratensis, 264  
 Podostroma, 179  
 Polyblastia rugulosa, 122, 123  
 Polychidium muscicolum, 138  
 Polycoccus punctiformis, 118, 138  
 Polyporus, 88, 313, 314; **albo-sordescens**, 88; **lucidus**, 115; **rufopodex**, 88  
 Polyscytalum, 45, 46, 55; **cylindroides**, 55; **fecundissimum**, 55; **sericeum**, 55, 56  
 Polystictus, 313  
 Populus fastigiata, 245; **grandiden-tata**, 238, 239  
 Porodisculus, 313  
 Poronidulus conchifer, 289  
 Probable identity of Stropharia epi-myces (Peck) Atk. with Pilosace algeriensis Fries, The, 167  
 Propolidium, 7; **Tsugae**, 10  
 Propolis faginea, 7  
 Protococcus, 127  
 Protoglossum, 313  
 Protuberia, 313  
 Prunus, 51  
 Psathyra, 36  
 Pseudoplectania, 299; **fulgens**, 299, 302; **melaena**, 300; **melania**, 299; **nigrella**, 299, 301, 302; **stygia**, 299, 301; **vogesiacae**, 299, 300, 302  
 Pseudoplectania, The genus, 299  
 Psidium, 19  
 Psilocybe uda, 314  
 Psora ostreata, 150  
 Psychotria glabrata, 27  
 Psychogaster, 314  
 Puccinia, 9; **angustata**, 303, 304; **caricina**, 240; **Caricis**, 240; **graminis**, 240, 304; **Peckiana**, 281; **striato-spora**, 71  
 Pucciniastrum Myrtilli, 237-239  
 Pulparia spongiosa, 300  
 Pycnochytrium, 315  
 Pycnosporos or ascospores, A method of determining in analytic work whether colonies of the chestnut blight fungus originate from, 274  
 Pyrenochaeta **fraxinina**, 247  
 Pyrenula nitida, 120, 124  
 Pyxidiphora, 176  
 Quercus, 58; **alba**, 58  
 Racoblenna, 119  
 Ramalina, 133; **calicaris**, 125  
 Ramularia, 180  
 Raphidium polymorphum, 131  
 Remarks on some Hypocreaceae, Maire's, 176  
 Reviews, News, notes and, 170  
 Rhinotrichum, 48; **Curtisii**, 48; **pul-veraceum**, 57  
 Rhizocarpon, 114; **alboatrum**, 114; **geographicum**, 114  
 Rhymbocarpus, 114; **punctiformis**, 114  
 Ribes, 242  
 Ricker, P. L., Ridgway's new color guide, 172  
 Ridgway's new color guide, 172  
 Rinodina sophodes, 127, 148  
 Rivularia, 119, 138; **nitida**, 118, 119  
 Roccella, 124, 140; **phycopsis**, 124  
 Rosa villosa, 247  
 Rostkovites granulatus, 3; **subaureus**, 3  
 Rubus, 245  
 Russula, 305, 308; **abietina**, 309; **albi-dula**, 310; **alutacea**, 310; **azurea**, 311; **bicolor**, 311; **brevipes**, 308; **chameleontina**, 309; **crenulata**,

- 310; *delica*, 308; *deliciosa*, 308; *drimeja*, 309; *emetica*, 76, 309; *flaviceps*, 310; *granulata*, 309; **Murrillii**, 310; *nigrescens*, 308; *nigricans*, 308; *obscura*, 309; *pectinata*, 311; *Turci*, 309; *veterinosa*, 309
- Rusts, Further cultures of heteroeccious, 233
- Salix*, 238, 239
- Sarcodon*, 12; *acre*, 13; *fuligineo-violaceus*, 13; **fumosus**, 16; *imbricatus*, 14; *laevigatus*, 14; **Murrillii**, 15; **radicatus**, 13; *reticulatus*, 12; **roseolus**, 16
- Sarcodon*, The genus. Type studies in the *Hydnaceae*—III, 12
- Sarcoscypha*, 185, 186; *coccinea*, 186; *Colensoi*, 191; *striispora*, 188
- Scenedesmus obtusus*, 131
- Scirpus*, 303
- Sclerodon*, 296
- Sclerotinia*, 109; *fructigena*, 50; *tuberosa*, 109
- Scutiger*, 290; *griseus*, 289
- Scytonema*, 113, 118, 124–126, 136, 138; *myochrous*, 109
- Seaver, F. J., Maire's remarks on some *Hypocreaceae*, 176; Shear's studies of parasitic species of *Glomerella*, 171; Some tropical cup-fungi, 185; The genus *Pseudoplectania*, 209; Wollenweber's studies on the *Fusarium* problem, 178
- Septoria*, 245; **Carriceriae**, 246
- Seurattia Coffeicola*, 40; *Vanillae*, 40
- Shear's studies of parasitic species of *Glomerella*, 171
- Simblum*, 267–269, 271, 273; *australe*, 265; *Lorentzii*, 265; *periphragmoides*, 266, 269; *pilidiatum*, 265; *rubescens*, 265; *sphaerocephalum*, 264, 266, 267, 269–272
- Simblum sphaerocephalum*, The structure of, 264
- Sirosiphon*, 116, 119, 124, 136, 138; *alpinus*, 109; *pulvinatus*, 109, 118
- Smith, C. S., Clark, E. D., & Toxicological studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*, 224
- Solidago*, 242
- Solorina crocea*, 118
- Some tropical cup-fungi, 185
- Species of *Clitocybe* and *Melanoleuca*, New. The *Agaricaceae* of the Pacific Coast—IV, 206
- Species of fungi from various localities, Notes on new, 245
- Species of fungi, New, 67
- Species of *Glomerella*, Shear's studies of parasitic, 171
- Sphaerella Opuntiae*, 38
- Sphaeria Lactifluorum*, 179
- Sphaeropsis Cocolobae*, 246; **rhodocarpa**, 246
- Sphaerotheca pannosa*, 58
- Sphagnum*, 300–302
- Spilonema paradoxum*, 119
- Sporendonema*, 45; *Casei*, 45; *myophilum*, 46; *terrestre*, 45
- Steccherinum*, 293
- Stereocaulon*, 118
- Sterility in *Pholiota candicans* (Bull.) Schroet., 314
- Stichococcus bacillaris*, 115, 131, 132; *bacillaris fungicola*, 115
- Stictis*, 7; *Tsugae*, 7, 10, 11
- Stigeoclonium tenue*, 131
- Stigmatomma cataleptum*, 121
- Stigonema mamillosum*, 109
- Strigula*, 120, 151; *complanata*, 120
- Stropharia*, 167, 168; *coprinophila*, 167, 168; *epimyces*, 168
- Stropharia epimyces* (Peck) Atk. with *Pilosace algeriensis* Fries, The probable identity of, 167
- Structure of *Simblum sphaerocephalum*, The, 264
- Studies in North American *Hyphomycetes*—II, 45
- Studies in the *Hydnaceae*, Type—III. The genus *Sarcodon*, 12; IV. The genus *Phellodon*, 62; V. The genus *Hydnellum*, 194; VI. The genera *Creolophus*, *Echinodontium*, *Gloiodon*, and *Hydnodon*, 293
- Studies of parasitic species of *Glomerella*, Shear's, 171
- Studies on the *Fusarium* problem, Wollenweber's, 178
- Studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*. Toxicological, 224
- Sumstine, D. R., Studies in North American *Hyphomycetes*—II, 45
- Synalissa*, 124
- Synchytrium*, 313; *Amsinckiae*, 313; *andinum*, 313; *aureum*, 315; *innominatum*, 313; *papillatum*, 313; *pluriannulatum*, 315; *Succisiae*, 315; *Taraxaci*, 315
- Synchytrium*, The genus, 315
- Taphrina*, 88; *lapponica*, 89
- Thelephora*, 105; *padinaeformis*, 297, 298
- Thelotrema lepadinum*, 122
- Thuja*, 6–8, 11; *occidentalis*, 10
- Thyridaria tarda*, 39
- Tilia*, 294

- Torula, 45, 53, 60; aurea, 48; fructigena, 49, 50; pallida, 60
- Toruloidea**, 53, 60; **candidula**, 53, 55, 61; **effusa**, 53, 61; **Nicotianae**, 53, 54; **Tulipiferae**, 53, 54, 61; **Unangstii**, 53, 54, 61
- Toxicological studies on the mushrooms *Clitocybe illudens* and *Inocybe infida*, 224
- Trametes*, 313; *Pini*, 115
- Trentepohlia*, 110, 116, 120, 124, 125, 127, 132, 136, 138-140, 147; aurea, 115, 116; umbrina, 109, 118, 119
- Trichoderma aureum*, 47
- Tricholoma*, 206, 208; album, 207, 220; **anomalum**, 223; **arenicola**, 223; **avellaneifolium**, 223; **avellaneum**, 223; **bicolor**, 223; **californicum**, 223; **collybiiforme**, 223; **dryophilum**, 223; **farinaceum**, 223; **Harperi**, 223; **luridum**, 221; **maculatescens**, 209; **nuciolens**, 223; **nudum**, 208; **Olesonii**, 223; **oreades**, 223; **pes-sundatum**, 222; **pinicola**, 223; **platyphyllum**, 223; **portolense**, 223; **roseibrunneum**, 223; **ru-dericolum**, 223; **secedifolium**, 223; **striatellum**, 223; **subluridum**, 223; **submulticeps**, 223; **subpessundatum**, 223; **subvelatum**, 223; **tenuipes**, 223
- Trichoscypha*, 185, 187; *Afzelii*, 189; *Hindsii*, 189; *insititia*, 190; *sul-cipes*, 189; *Tricholoma*, 188
- Tropical cup-fungi, Some, 185
- Tropical North America, The Agari-caceae of—IV, 18
- Tsuga*, 6-8, 11, 233; *canadensis*, 10, 237-239
- Type studies in the Hydnaceae—III. The genus *Sarcodon*, 12; IV. The genus *Phellodon*, 62; V. The genus *Hydnellum*, 194; VI. The genera *Creolophus*, *Echinodontium*, *Gloio-don*, and *Hydnodon*, 293
- Ulothrix*, 125
- Uncinula necator*, 58
- Uredinales on *Carex* in North Amer-ica, 240
- Uredinopsis, 223, 237, 304; *Atkin-sonii*, 236, 239; *mirabilis*, 236, 239; *Osmundae*, 235, 239; *Phegopteridis*, 236, 239; *Struthiopteridis*, 234, 235, 239
- Uredo Scabies, 40
- Uromyces Caladii*, 304
- Urophlyctis pluriannulatus*, 313
- Urtica*, 242
- Vaccinium canadense*, 237-239; *penn-sylvanicum*, 52, 238
- Vaginata*, 81, 94; *agglutinata*, 82-85, 94; **albocreata**, 79, 82, 84; *fari-nosa*, 82-85; *livida*, 82; **parcivol-vata**, 82, 83; **plumbea**, 82, 83, 85, 86; *plumbea alba*, 84; *plumbea strangulata*, 82; **pusilla**, 82, 83; *vaginata*, 82
- Valsaria Akebiae*, 248
- Vaucheria sessilis*, 127
- Venenarius*, 72, 81, 94, 258; **Caesar-eus**, 72, 73, 85; **crenulatus**, 73, 77; **cothurnatus**, 72, 74, 77, 79, 81, 96; **flavorubescens**, 73, 76, 95; **Frosterianus**, 73, 76, 79, 96; **junquilleus**, 80; **Morrisii**, 73, 75; *muscarius*, 73, 75-77, 94, 96; **pantherinus**, 80, 81, 84; *phal-loides*, 72, 74, 79-81, 93; **por-phyrius**, 81; **recutitus**, 81; **rubens**, 73, 75, 76, 79, 95; **rus-suloides**, 73, 76, 77, 80; *solitarius*, 73, 77, 79, 85; **spissus**, 81; **spretus**, 72, 73, 81; **velatipes**, 73, 75
- Verrucaria*, 124; *muralis*, 123, 127
- Verticillium*, 180
- Wilson, G. W., The genus *Synchy-trium*, 315
- Wolf, F. A., Internal aecia, 303
- Wollenweber's studies on the Fu-sarium Problem, 178
- Wynnea, 186
- Xanthoria*, 128; *parietina*, 125, 127-132, 151
- Xylaria*, 178
- Zea Mays*, 52
- Zygodesmus*, 60